



Prevalence of carbapenem resistance genes in Enterobacteriaceae, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* from meat samples

Maryam Rezaeian¹ , Laleh Hoveida^{1*}

1. Department of Microbiology, Fal.C., Islamic Azad University, Isfahan, Iran

* Correspondence: Laleh Hoveida, Department of Microbiology, Fal.C., Islamic Azad University, Isfahan, Iran.

Tel: +989121480576; Email: La.hoveida@iau.ac

Article Type: Research Article

Article History

Received: 10 September 2024

Received in revised form: 15 October 2024

Accepted: 29 December 2024

Available online: 22 February 2025

DOI: [10.29252/mlj.19.1.36](https://doi.org/10.29252/mlj.19.1.36)

Keywords

[Enterobacteriaceae](#)
[Acinetobacter baumannii](#)
[Pseudomonas aeruginosa](#)
[Carbapenems](#)

Abstract

Background: Infections caused by bacteria transmitted from food, including carbapenem-resistant *Enterobacteriaceae*, *Pseudomonas aeruginosa* (*P. aeruginosa*), and *Acinetobacter baumannii* (*A. baumannii*), are spreading, and this has caused concerns in treatment. This study investigated the frequency of carbapenem resistance genes in *Enterobacteriaceae*, *P. aeruginosa*, and *A. baumannii*, isolated from raw chicken meat.

Methods: In this cross-sectional study, 100 samples of raw chicken meat were collected from Isfahan. Bacterial contamination was evaluated and confirmed using biomedical tests. Antibiotic sensitivity tests were performed using disc diffusion for *Enterobacteriaceae*, *P. aeruginosa*, and *Acinetobacter baumannii* isolates. The frequency of carbapenem resistance genes (*OXA-181*, *OXA-48*, *VIM*, *NDM*, *IMP*, and *KPC*) was determined through polymerase chain reaction (PCR) analysis.

Results: Out of 100 samples, 70 were positive for bacterial infection, of which 30 were infected with *Escherichia coli* (*E. coli*), 14 samples with *Klebsiella pneumoniae* (*K. pneumoniae*), 9 samples with *Salmonella typhimurium* (*S. typhimurium*), 11 samples with *P. aeruginosa*, and 6 samples with *A. baumannii*. The highest levels of antibiotic resistance were estimated to be tetracycline, cotrimoxazole, gentamicin, trimethoprim, and streptomycin, and the lowest levels were azithromycin and rifampin. Among genes encoding resistance to carbapenem, *NDM* and *OXA-48* genes were the most commonly expressed, with a frequency of 60% and 28.24%, respectively.

Conclusion: This study found significant bacterial contamination, especially for *Enterobacteriaceae*, with notable antibiotic resistance to tetracycline. Carbapenem resistance genes *NDM* and *OXA-48* were prevalent, indicating the urgency of addressing antibiotic resistance.



OPEN ACCESS



© The author(s)

Introduction

As the global population expands, ensuring food security becomes increasingly pressing. It is necessary to assess whether food production systems can meet this demand while maintaining high quality and adhering to safety protocols. Animal-derived food products, such as meat, are an important part of the human diet, yet their safety and sustainability are threatened by various factors (1,2). Foodborne infections caused by Gram-positive and Gram-negative bacteria are becoming more widespread, raising concerns about treatment efficacy (3). The prevalence of antibiotic resistance in the food chain poses a significant threat to human health, as antibiotic-resistant bacteria can enter the food supply through various pathways, including the use of antibiotics in livestock for growth promotion and disease prevention (4,5). Among the diverse array of resistance mechanisms, the rise of carbapenem resistance in clinically relevant pathogens represents a pressing concern due to the limited treatment options available (6,7). The mechanisms of antibiotic resistance -such as the acquisition of resistance genes and the production of carbapenemase enzymes that render carbapenems ineffective- highlight the complexity of this issue, as these enzymes can spread among bacterial species, complicating treatment options (8,9).

Carbapenems, including imipenem and meropenem, constitute crucial medications renowned for their extensive and enduring spectrum of activity in contrast to beta-lactamases (10). *Enterobacteriaceae*, *P. aeruginosa*, and *A. baumannii* are notable bacterial species known for acquiring and disseminating resistance genes, including those conferring resistance to carbapenems (8,11-14). The most important carbapenems are commonly categorized into three classes: Class A (*KPC*), Class B: encompassing metallo β -lactamases such as *IMP*, *VIM*,

and *NDM*, and Class D: represented by oxacillinase, notably *OXA-48*, of which *OXA-48* has been reported frequently worldwide for its role in the rapid spread of infection (15,16).

Meatborne diseases caused by bacterial infections are more common in children (17). Numerous bacterial pathogens, including *P. aeruginosa*, *A. baumannii*, *Klebsiella pneumoniae* (*K. pneumoniae*), *E. coli*, *Salmonella*, *Listeria monocytogenes*, *Yersinia enterocolitica*, toxin-producing species like *Staphylococcus aureus* (*S. aureus*), and *Bacillus cereus*, contribute to meatborne diseases either through animal infection or contamination during meat processing or handling (18-20). Consuming contaminated meat poses a risk of contracting various diseases, categorized into gastrointestinal and extra-gastrointestinal infections (21). In the current study, we investigated the frequency of carbapenem resistance genes in *Enterobacteriaceae*, *P. aeruginosa*, and *A. baumannii* isolated from raw chicken meat in Isfahan, Iran.

Methods

One hundred raw chicken meat samples were collected from various farms to represent different production settings and potential contamination levels. The samples were obtained using sterile swabs to prevent any external contamination during collection. Each sample was cultured on Eosin Methylene Blue (EMB) medium, adhering to standard microbiological protocols to promote the growth of Gram-negative bacteria. Following the initial confirmation of Gram-negative bacilli, positive lactose colonies were further cultured on a Triple Sugar Iron Agar medium specifically designed for the isolation of *Enterobacteriaceae*. For the identification of *Pseudomonas* species, suspicious colonies were cultured on *Pseudomonas* Cefrimide Agar (PCA), while colonies suspected to be *A. baumannii* were cultured on

blood agar. Phenotypic and biochemical tests were applied to identify the isolates as follows: for *Enterobacteriaceae*, indole, methyl red, Voges Proskauer, and citrate tests were utilized to differentiate diagnosis between *E. coli*, *S. typhimurium*, and *K. pneumoniae*; *P. aeruginosa* isolated were confirmed through lactose, citrate, indole, oxidase, DNase, and hemolysis tests; and *A. baumannii* isolates were confirmed by lactose, oxidase, catalase, pigmentation, urease, and IMVIC tests.

The antibiotic susceptibility of all isolates was determined by the Kirby-Bauer method. Applied antibiotics included tetracycline, ceftazidime, ciprofloxacin, trimethoprim-sulfamethoxazole, tobramycin, chloramphenicol, norfloxacin, amikacin, gentamicin, rifampin, cephalothin, streptomycin, trimethoprim, levofloxacin, imipenem, meropenem, azithromycin (22).

DNA extraction was conducted from pure bacterial cultures (23) and was assessed using NanoDrop (24). Polymerase chain reaction (PCR) to detect *OXA-181*, *OXA-48*, *VIM*, *NDM*, *IMP* and *KPC* genes expression was performed through the following steps: for *E. coli* (1 for 5 min cycle at 94 °C for initial denaturation; 33 cycles included 60 sec at 94 °C for denaturation, 45 sec at 58 °C for annealing, and 60 sec at 72 °C for extension; and 1 cycle for 7 min at 72 °C for final extension), for *K. pneumoniae* (1 for 5 min cycle at 94 °C for initial denaturation; 35 cycles included 30 sec at 95 °C for denaturation, 90 sec at 58 °C for annealing, and 90 sec at 72 °C for extension; and 1 cycle for 10 min at 72 °C for final extension), for *S. typhimurium* (1 for 4 min cycle at 95 °C for initial denaturation; 30 cycles included 45 sec at 94 °C for denaturation, 60 sec at 58 °C for annealing, and 40 sec at 72 °C for extension; and 1 cycle for 5 min at 72 °C for final extension), for *P. aeruginosa* (1 for 5 min cycle at 94 °C for initial denaturation; 25 cycles included 35 sec at 94 °C for denaturation, 45 sec at 53 °C for annealing, and 60 sec at 72 °C for extension; and 1 cycle for 7 min at 72 °C for final extension), for *A. baumannii* (1 for 3 min cycle at 94 °C for initial denaturation; 30 cycles included 40 sec at 95 °C for denaturation, 55 sec at 59 °C for annealing, and 60 sec at 72 °C for extension; and 1 cycle for 6 min at 72 °C for final extension). Multiplex PCR was performed to evaluate the expression of gene coding for resistance to carbapenem compounds in isolates. Utilized primers have been presented in Table 1 (25). The quality of the product was evaluated using gel electrophoresis (26).

Statistical analysis

Numbers and percentages reported the frequency of data. All data was analyzed using SPSS version 21.

Table 1. Forward and reverse primers

Genes/Primers name	Sequence (5'-3')	Product size (bp)
<i>16sr RNA</i> (<i>E. coli</i>)	F:AGATTGATCMTGGCTCAG R:CCGTC AATTCATTTGATTT	919
<i>16S-23SITS</i> (<i>K. pneumoniae</i>)	F:ATTTGAAGAGGTTGCAAACGAT R:TTCACTCTGAAGTTTCTGTGTTC	130
<i>23sr RNA</i> (<i>S. typhimurium</i>)	F:GCCAACCATTGCTAAATTGGCGCA R:GGTAGAAATCCAGCGGGTACTGG	429
<i>NanI</i> (<i>P. aeruginosa</i>)	F:ATGAATACTTATTTTGATAT R:CTAAATCCATGCTCTGACCC	228
<i>16s-23srDNA</i> (<i>A. baumannii</i>)	F:CATTATCACGGTAATTAGTG R:AGAGCACTGTGCACTTAAG	208
<i>KPC</i>	F:ATTTTCAGAGCCTTACTGCCC R:TATCGTTGATGCTACTGTATCG	901
<i>IMP</i>	F:GGAATAGAGTGGCTTAAYTCTC R:GGTTAAAYAAAAACAACCACC	232
<i>VIM</i>	F:GATGGTGTGGTGCATA R:CGAATGCGCAGCACCAG	390
<i>NDM</i>	F:CCGTATGAGTGATTGCGGGC R:GCCCAATATTATGCAACCCGG	779
<i>OXA-48 & OXA181</i>	F:TATATTGCATTAGCAAGGG R:CACACAATACGCGCTAACC	847

Results

Studied population and bacterial isolates

Among 100 cultured samples, 70 were positive for bacterial infections. Following standard bacteriological tests, 30 isolates were identified as *E. coli*, 14 isolates as *K. pneumoniae*, 9 isolates as *S. typhimurium*, 11 as *P. aeruginosa*, and 6 isolates were identified as *A. baumannii*.

Antibiotic susceptibility test

The highest rates of antibiotic resistance were found in *E. coli* against tetracycline (100%) and streptomycin (93.33%), while *K. pneumoniae* also showed 100% resistance to tetracycline. *S. typhimurium* was resistant to tetracycline (100%) and trimethoprim (88.88%). *P. aeruginosa* exhibited 100% resistance to tetracycline, and *A. baumannii* showed complete resistance (100%) to meropenem and imipenem. Notably, *P. aeruginosa* showed the highest resistance to imipenem (Table 2, Figure 1).

Table 2. Antibiotic susceptibility results

Antibiotic	Resistance n (%)				
	<i>Escherichia coli</i> (n=30)	<i>Klebsiella pneumoniae</i> (n=14)	<i>Salmonella typhimurium</i> (n=9)	<i>Pseudomonas aeruginosa</i> (n=11)	<i>Acinetobacter baumannii</i> (n=6)
Streptomycin	28 (93.33)	12 (85.71)	6 (66.67)	3 (27.27)	3 (50.00)
Gentamicin	22 (73.33)	10 (71.43)	8 (88.89)	10 (90.91)	6 (100.00)
Ampicillin	14 (46.67)	8 (57.14)	4 (44.44)	5 (45.45)	5 (83.33)
Tobramycin	8 (26.67)	5 (35.71)	2 (2.22)	6 (54.55)	4 (66.67)
Cotrimoxazole	26 (86.67)	10 (71.43)	8 (88.89)	9 (81.82)	3 (50.00)
Cephalothin	14 (46.67)	5 (35.71)	0 (0.00)	8 (72.73)	3 (50.00)
Ceftazidime	16 (53.33)	6 (42.86)	0 (0.00)	9 (81.82)	3 (50.00)
Tetracycline	30 (100)	14 (100)	9 (100)	11 (100)	6 (100)
Trimethoprim	26 (86.67)	10 (71.43)	8 (88.89)	8 (72.73)	3 (50.00)
Ciprofloxacin	8 (26.67)	6 (42.86)	6 (66.67)	6 (54.55)	5 (83.33)
Levofloxacin	9 (30.00)	2 (14.29)	2 (22.22)	8 (72.73)	5 (83.33)
Imipenem	8 (26.67)	2 (14.29)	0 (0.00)	9 (81.82)	6 (100)
Meropenem	7 (23.33)	3 (21.43)	0 (0.00)	6 (54.55)	6 (100)
Chloramphenicol	11 (36.67)	8 (57.14)	6 (66.67)	2 (18.18)	2 (33.33)
Nitrofurantoin	9 (30.00)	7 (50.00)	8 (88.89)	0 (0.00)	2 (33.33)
Azithromycin	7 (23.33)	2 (14.29)	0 (0.00)	8 (72.73)	4 (66.67)
Rifampin	8 (26.67)	0 (0.00)	2 (22.22)	0 (0.00)	3 (50.00)

Molecular detection of carbapenemase genes

The most frequently reported gene expression of *E. coli*, *K. pneumoniae*, and *S. typhimurium* was *NDM* (7.33%), *KPC* (71.43%), and *NDM* (66.67%), respectively, which were significantly higher than those of other genes ($p < 0.05$). Among *P. aeruginosa* isolates, the frequency of *VIM* (100%) and *IMP* (90.91%) gene expression was significantly higher compared to others ($p = 0.018$). In *A. baumannii* isolates, the gene

expression of *IMP*, *VIM*, and *OXA-48* was significantly higher than others ($p = 0.030$). Notably, the presence of the *NDM* gene in *S. typhimurium*, which did not show carbapenem resistance, highlighted the complexity of antibiotic resistance gene distribution, suggesting that resistance genes should primarily be reported in the context of resistant strains to provide clearer implications for resistance mechanisms (Table 3, Figure 2).

Table 3. Frequency of carbapenemase gene expression among bacterial isolates

Bacterial isolates	Frequency of gene expression n (%)						P-value
	<i>KPC</i>	<i>IMP</i>	<i>VIM</i>	<i>NDM</i>	<i>OXA-48</i>	<i>OXA-181</i>	
<i>Escherichia coli</i> (n=30)	14 (46.67)	16 (53.33)	14 (46.67)	22 (73.33)	6 (20.00)	16 (53.33)	0.023
<i>Klebsiella pneumoniae</i> (n=14)	10 (71.43)	8 (7.14)	6 (42.86)	8 (57.14)	3 (21.43)	6 (42.86)	0.041
<i>Salmonella typhimurium</i> (n=9)	3 (33.33)	2 (22.22)	2 (22.22)	6 (66.67)	0 (0.00)	4 (44.44)	0.018
<i>Pseudomonas aeruginosa</i> (n=11)	6 (54.5)	10 (90.91)	11 (100)	4 (36.36)	2 (18.18)	3 (27.27)	0.044
<i>Acinetobacter baumannii</i> (n=6)	3 (50.00)	6 (100)	6 (100)	2 (33.33)	6 (100)	2 (33.33)	0.030




Figure 1. Resistance to carbapenem




Figure 2. Molecular detection of the most prevalent carbapenemase genes: A: *NDM* and B: *OXA48* and *OXA181*.

Discussion

The emergence and spread of antibiotic resistance among bacterial pathogens pose significant challenges to public health worldwide. The increasing prevalence of carbapenem resistance genes in various bacterial species has raised concerns about the potential transmission of these resistance genes through the food chain (6,7). The expression of carbapenemase genes, which facilitates bacterial survival against carbapenems, raises concerns about the effectiveness of current treatment protocols and the need for novel therapeutic strategies. Strengthening surveillance and implementing targeted infection control measures are essential to mitigate the impact of these resistant strains on public health (27,28). In the current study, we investigated the frequency of carbapenem resistance genes in Enterobacteriaceae, *P. aeruginosa*, and *A. baumannii* isolated from meat. Among Enterobacteria, 42.8% of isolates were *E. coli*, 20% were *K. pneumoniae*, and 12.9% were *S. typhimurium*. The frequency of *P. aeruginosa* was 15.7%, and for *A. baumannii* was 8.6%.

Similarly, Dehkordi et al. reported a prevalence of 45.2% for *E. coli* isolated from raw chicken meat in Iran (29). Nazari Moghadam et al. demonstrated that 2% of chicken meat samples were positive for *Salmonella*, and all were *S. typhimurium* (30). It indicates that most antibiotics commonly utilized in the livestock and poultry industries are ineffective against a significant isolate of *Salmonella*, underscoring their importance in public health considerations. In a study by Mousse et al., about 20% of bacteria isolated from the food were *K. pneumoniae* (31). Atabay et al. observed that among 484 isolates, 78.92% were *E. coli*, followed by 7.64% *Salmonella*, 3.71% *Proteus*, 3.51% *S. aureus*, 1.85% *Streptococcus*, 1.44% *P. aeruginosa*, 1.23% *Klebsiella*, and 1.03% *S. epidermidis* (32). Rafei et al. reported different prevalences of *A. baumannii* isolated from meat, raw milk, drinking water, cheese, and domestic animal samples, respectively (33).

We observed that about 70% of selected samples contained Gram-negative bacteria from raw chicken meat with high antibiotic resistance to various antibiotics, including tetracycline, cotrimoxazole, gentamicin, trimethoprim, and streptomycin. This high contamination rate likely results from the overuse of antibiotics in poultry farming, where these substances are routinely prescribed to control and prevent bacterial infections, treat diseases, and promote growth (1,34). Such practices not only contribute to the development of resistant bacterial strains but also pose significant public health risks as these resistant pathogens can be transmitted to humans through the food chain. Moreover, the widespread resistance observed may compromise the effectiveness of treatment options for bacterial infections in humans, necessitating urgent interventions to regulate antibiotic use in livestock and improve food safety standards (35).

Rezaloo et al. found that 9.16% of meat samples were *P. aeruginosa*. These isolates exhibited pronounced antibiotic resistance, particularly to ampicillin, penicillin, and tetracycline, and the lowest resistance was reported to imipenem and trimethoprim (36). Studies reported that *K. pneumoniae* and *E. coli* isolated from meat were more likely to be multidrug-resistant and resistant (37,38). *K. pneumoniae* isolated from meat was more resistant to tetracycline and gentamicin than those isolated from the gastrointestinal tract (37). Among detected *E. coli*, *K. pneumoniae*, *S. typhimurium*, and *P. aeruginosa* had the highest resistance to tetracycline. Yang et al. reported that among *Salmonella* isolated from meat, *S. typhimurium* isolates were more prevalent than others with the highest resistance to tetracycline (39). Consequently, the prescription of antibiotics and, inevitably, the emergence of antibiotic resistance in poultry farms, is exacerbated. The primary source of contamination in livestock and poultry stems from the transmission of antibiotic-resistant bacterial strains from humans to meat, leading to an escalation in the prevalence of strains isolated from livestock and poultry that are resistant to antibiotics used in human medicine (40,41).

Alhazmi et al.'s study on *K. pneumoniae* isolates showed a high frequency of *blaOXA-48* and *blaNDM* carbapenemase genes. Among their isolates, only one isolate each harbored *blaVIM* and *blaKPC* genes, while no *blaIMP*-producing isolates were detected (42). In the current study, KPC was the most prevalent carbapenemase gene expression in *K. pneumoniae*, while *OXA-48* had the lowest frequency. It should be considered that differences in geographical location and sample origin between studies may influence the variations of the genes. Bakhtiari et

al. demonstrated that 24% of *K. pneumoniae* isolates were resistant to imipenem, 35.45% represented the *blaKPC* gene, and 16.36% expressed the *blaVIM-1* gene (43). Amini et al. demonstrated that among *K. pneumoniae* isolates, expression of *blaVIM* and *blaIMP* were higher than other carbapenemase gene (44). Li et al. showed that among carbapenem-resistant *A. baumannii*, common carbapenemase-positive genes included *blaOXA-51*-like, *blaOXA-23*-like, *blaNDM-1*, and *blaOXA-58* (45). The current study reported gene expressions of *IMP*, *VIM*, and *OXA-48* genes in *A. baumannii* isolates. Another study by Ghaffoori Kanaan et al. demonstrated that the most common gene expression among carbapenem-resistant *S. enteritidis* isolates was *blaIMP*, *blaOXA-48*-like, and *blaNDM*. In contrast, the *blaKPC* and *blaVIM* genes were undetected (46). We detected *S. typhimurium* and *E. coli* from collected meat, with the highest frequency of *NDM* expression.

The differences observed in the levels of antibiotic resistance and the prevalence of specific genes across various studies are multifactorial. These include the indiscriminate use of antibiotics in different countries, variations in patterns of antibiotic administration, the emergence of novel resistance mechanisms, the site of infection acquisition, genetic variations among bacterial strains, and geographical conditions, among others (47,48). The emergence of antibiotic resistance poses a significant burden on healthcare systems globally, leading to substantial economic costs. Apart from inappropriate antibiotic usage, factors such as aggressive treatment approaches, immunocompromised patients, and non-compliance with hygiene protocols also contribute to the problem. In the agricultural sector, adherence to health regulations, proper animal nutrition, modern slaughtering practices, and maintaining the integrity of the cold chain during transportation play pivotal roles in infection control efforts.

This study has some limitations that should be acknowledged. First, the sample size was relatively small, which may affect the generalizability of our findings. Additionally, the geographical bias inherent in our sampling could limit the applicability of the results to broader contexts, as antibiotic resistance patterns may vary significantly across different regions due to variations in antibiotic usage and regulations. Finally, the potential for changes in resistance patterns over time, influenced by evolving antibiotic practices, cannot be ruled out. Future research with larger, more diverse samples is essential to provide a more comprehensive understanding of carbapenem resistance in these pathogens.

Conclusion

The current study underscored multifaceted factors influencing antibiotic resistance disparities among studies, including varied antibiotic usage patterns, emergence of resistance mechanisms, and genetic diversity among bacterial strains. Antibiotic resistance poses a substantial global healthcare burden, exacerbated by inappropriate usage and other factors like aggressive treatments and agricultural practices. Addressing these complexities is critical for effective infection control.

Acknowledgement

Not applicable.

Funding sources

None.

Ethical statement

The study was approved by the Ethics Committee of the Falavarjan Branch, Islamic Azad University, Isfahan, Iran (IR.IAU.FALA.REC.1400.033).

Conflicts of interest

The authors declare that they have no competing interests.

Author contributions

All authors made significant contributions to the work reported. MR and LH participated in the research design and writing the first draft; MR and LH performed the research and analysis tools; MR and LH participated in data analysis. All authors reviewed and confirmed the final manuscript.

Data availability statement

All data generated or analyzed during this study are included in this published article.

References

- Abreu R, Semedo-Lemsaddek T, Cunha E, Tavares L, Oliveira M. Antimicrobial drug resistance in poultry production: Current status and innovative strategies for bacterial control. *Microorganisms*. 2023;11:953. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Leroy F, Smith NW, Adesogan AT, Beal T, Iannotti L, Moughan PJ, et al. The role of meat in the human diet: evolutionary aspects and nutritional value. *Anim Front Rev Mag Anim Agric*. 2023;13:11-8. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Antunes P, Novais C, Peixe L. Food-to-Humans Bacterial Transmission. *Microbiol Spectr*. 2020;8(1):10.1128. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Kumar SB, Arnipalli SR, Ziouzenkova O. Antibiotics in Food Chain: The Consequences for Antibiotic Resistance. *Antibiot (Basel, Switzerland)*. 2020;9(10):688. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Manyi-Loh C, Mamphweli S, Meyer E, Okoh A. Antibiotic Use in Agriculture and Its Consequential Resistance in Environmental Sources: Potential Public Health Implications. *Molecules*. 2018;23(4):795. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Mancuso G, De Gaetano S, Midiri A, Zummo S, Biondo C. The Challenge of Overcoming Antibiotic Resistance in Carbapenem-Resistant Gram-Negative Bacteria: "Attack on Titan". *Microorganisms*. 2023;11(8):1912. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Salam MA, Al-Amin MY, Salam MT, Pawar JS, Akhter N, Rabaan AA, et al. Antimicrobial resistance: a growing serious threat for global public health. In: *Healthcare*. MDPI. 2023;1946. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Mancuso G, Midiri A, Gerace E, Biondo C. Bacterial Antibiotic Resistance: The Most Critical Pathogens. *Pathog (Basel, Switzerland)*. 2021;10. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Muteeb G, Rehman MT, Shahwan M, Aatif M. Origin of Antibiotics and Antibiotic Resistance, and Their Impacts on Drug Development: A Narrative Review. *Pharmaceuticals (Basel)*. 2023;16(11):1615. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Sawa T, Kooguchi K, Moriyama K. Molecular diversity of extended-spectrum β -lactamases and carbapenemases, and antimicrobial resistance. *J Intensive Care*. 2020;8:13. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Kyriakidis I, Vasileiou E, Pana ZD, Tragiannidis A. *Acinetobacter baumannii* Antibiotic Resistance Mechanisms. *Pathog (Basel, Switzerland)*. 2021;10(3):373. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Sheu C-C, Chang Y-T, Lin S-Y, Chen Y-H, Hsueh P-R. Infections Caused by Carbapenem-Resistant Enterobacteriaceae: An Update on Therapeutic Options. *Front Microbiol*. 2019;10:80. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Hasannejad-Bibalan M, Sadeghi M, Hemmati H, Ashoobi MT, Yaghoubi T, Samadnia A, et al. A two-year study of microbiological characteristics of intravascular catheter-related bloodstream infections at Razi hospital, Iran. *New Zeal J Med Lab Sci*. 2021;75:202-5. [View at Publisher] [Google Scholar]
- Hosseini M, Hasannejad-Bibalan M, Yaghoubi T, Mobayen M, Khoshdoz P, Khoshdoz S, et al. Prevalence and Antibiotic Resistance Pattern of Gram-Positive Isolates from Burn Patients in Velayat Burn Center in Rasht, North of Iran. *Med Lab J*. 2021;15:52-7. [View at Publisher] [DOI] [Google Scholar]
- Saleh A, Göttig S, Hamprecht AG. Multiplex Immunochromatographic Detection of OXA-48, KPC, and NDM Carbapenemases: Impact of Inoculum, Antibiotics, and Agar. *J Clin Microbiol*. 2018;56(5):e00050-18. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Pitout JDD, Peirano G, Kock MM, Strydom KA, Matsumura Y. The Global Ascendancy of OXA-48-Type Carbapenemases. *Clin Microbiol Rev*. 2019;33(1):e00102-19. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Heredia N, García S. Animals as sources of food-borne pathogens: A review. *Anim Nutr (Zhongguo xu mu shou yi xue hui)*. 2018;4:250-5. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Bintsis T. Foodborne pathogens. *AIMS Microbiol*. 2017;3:529-63. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Li X, Gu N, Huang TY, Zhong F, Peng G. *Pseudomonas aeruginosa*: A typical biofilm forming pathogen and an emerging but underestimated pathogen in food processing. *Front Microbiol*. 2022;13:1114199. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Elbehiry A, Marzouk E, Moussa IM, Dawoud TM, Mubarak AS, Al-Sarar D, et al. *Acinetobacter baumannii* as a community foodborne pathogen: Peptide mass fingerprinting analysis, genotypic of biofilm formation and phenotypic pattern of antimicrobial resistance. *Saudi J Biol Sci*. 2021;28:1158-66. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Ali S, Alsayeqh AF. Review of major meat-borne zoonotic bacterial pathogens. *Front public Heal*. 2022;10:1045599. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Rai S, Dash D, Agarwal N. Introducing the new face of CLSI M100 in 2023: An explanatory review. *Indian J Med Microbiol*. 2023;46:100432. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Ghiaei A, Ghasemi SM, Shokri D. Investigating the antagonistic effect of indigenous probiotics on carbapenem-resistant *Pseudomonas aeruginosa* strains. *Biomed Res Int*. 2023;2023. [View at Publisher] [DOI] [PMID] [Google Scholar]
- García-Alegria AM, Anduro-Corona I, Pérez-Martínez CJ, Guadalupe Corella-Madueño MA, Rascón-Durán ML, Astiazaran-García H. Quantification of DNA through the NanoDrop Spectrophotometer: Methodological Validation Using Standard Reference Material and Sprague Dawley Rat and Human DNA. *Int J Anal Chem*. 2020;2020:8896738. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Shoja S, Ansari M, Faridi F, Azad M, Davoodian P, Javadpour S, et al. Identification of carbapenem-resistant *Klebsiella pneumoniae* with emphasis on New Delhi metallo-beta-lactamase-1 (bla NDM-1) in Bandar Abbas, South of Iran. *Microb Drug Resist*. 2018;24:447-54. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Ziraldo R, Shoura MJ, Fire AZ, Levene SD. Deconvolution of nucleic-acid length distributions: a gel electrophoresis analysis tool and applications. *Nucleic Acids Res*. 2019;47:e92. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Caliskan-Aydogan O, Alocilja EC. A Review of Carbapenem Resistance in Enterobacteriales and Its Detection Techniques. *Microorganisms*. 2023;11. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Bonomo RA, Burd EM, Conly J, Limbago BM, Poirel L, Segre JA, et al. Carbapenemase-Producing Organisms: A Global Scourge. *Clin Infect Dis an Off Publ Infect Dis Soc Am*. 2018;66:1290-7. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Dehkordi FS, Yahaghi E, Darian EK. Prevalence of antibiotic resistance in *Escherichia coli* isolated from poultry meat supply in Isfahan. *Iran J Med Microbiol*. 2014;8(2):41-7 [View at Publisher] [Google Scholar]
- Nazari Moghadam M, Rahimi E, Shakerian A, Momtaz H. Prevalence of *Salmonella Typhimurium* and *Salmonella Enteritidis* isolated from poultry meat: virulence and antimicrobial-resistant genes. *BMC Microbiol*. 2023;23:168. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Moussé W, Noumavo PA, Chabi NW, Sina H, Tohoeyessou MG, Ahoyo TA, et al. Phenotypic and genotypic characterization of extended spectrum β -Lactamase *Klebsiella pneumoniae* and fluorescent *Pseudomonas* spp. strains from market garden products and their watering water in Benin (West Africa). *Food Nutr Sci*. 2016;7:192-204. [View at Publisher] [DOI] [Google Scholar]

32. Atabay Z, Peighambari SM, Madani SA, Yazdani A. Bacteriologic survey of hepatic and cardiac lesions in commercial poultry carcasses. *Iranian Veterinary Journal*; 2023;18(4):5-14. [[View at Publisher](#)] [[DOI](#)] [[Google Scholar](#)]
33. Rafei R, Hamze M, Pailhoriès H, Eveillard M, Marsollier L, Joly-Guillou M-L, et al. Extrahuman epidemiology of *Acinetobacter baumannii* in Lebanon. *Appl Environ Microbiol*. 2015;81:2359-67. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
34. Roth N, Käsbohrer A, Mayrhofer S, Zitz U, Hofacre C, Domig KJ. The application of antibiotics in broiler production and the resulting antibiotic resistance in *Escherichia coli*: A global overview. *Poult Sci*. 2019;98:1791-804. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
35. Odey TOJ, Tanimowo WO, Afolabi KO, Jahid IK, Reuben RC. Antimicrobial use and resistance in food animal production: food safety and associated concerns in Sub-Saharan Africa. *Int Microbiol*. 2024;27:1-23. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
36. Rezaloo M, Motalebi A, Mashak Z, Anvar A. Prevalence, antimicrobial resistance, and molecular description of *Pseudomonas aeruginosa* isolated from meat and meat products. *J Food Qual*. 2022;2022:1-11. [[View at Publisher](#)] [[DOI](#)] [[Google Scholar](#)]
37. Davis GS, Waits K, Nordstrom L, Weaver B, Aziz M, Gauld L, et al. Intermingled *Klebsiella pneumoniae* Populations Between Retail Meats and Human Urinary Tract Infections. *Clin Infect Dis*. 2015;61:892-9. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
38. Rahman MM, Husna A, Elshabrawy HA, Alam J, Runa NY, Badruzzaman ATM, et al. Isolation and molecular characterization of multidrug-resistant *Escherichia coli* from chicken meat. *Sci Rep*. 2020;10:21999. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
39. Yang X, Wu Q, Zhang J, Huang J, Chen L, Wu S, et al. Prevalence, Bacterial Load, and Antimicrobial Resistance of *Salmonella* Serovars Isolated from Retail Meat and Meat Products in China. *Front Microbiol*. 2019;10:2121. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
40. Kousar S, Rehman N, Javed A, Hussain A, Naem M, Masood S, et al. Intensive Poultry Farming Practices Influence Antibiotic Resistance Profiles in *Pseudomonas aeruginosa* Inhabiting Nearby Soils. *Infect Drug Resist*. 2021;14:4511-6. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
41. Mtonga S, Nyirenda SS, Mulemba SS, Ziba MW, Muuka GM, Fandamu P. Epidemiology and antimicrobial resistance of pathogenic *E. coli* in chickens from selected poultry farms in Zambia. *J Zoonotic Dis*. 2021;5:18-28. [[View at Publisher](#)] [[Google Scholar](#)]
42. Alhazmi W, Al-Jabri A, Al-Zahrani I. The Molecular Characterization of Nosocomial Carbapenem-Resistant *Klebsiella pneumoniae* Co-Harboring bla NDM and bla OXA-48 in Jeddah. *Microbiol Res (Pavia)*. 2022;13:753-64. [[View at Publisher](#)] [[DOI](#)] [[Google Scholar](#)]
43. Bakhtiari R, Javadi A, Aminzadeh M, Molaei-Aghaei E, Shaffaghat Z. Association between Presence of RmpA, MrkA and MrkD Genes and Antibiotic Resistance in Clinical *Klebsiella pneumoniae* Isolates from Hospitals in Tehran, Iran. *Iran J Public Health*. 2021;50:1009-16. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
44. Amin M, Navidifar T, Saleh Shooshtari F, Goodarzi H. Association of the genes encoding metallo- β -lactamase with the presence of integrons among multidrug-resistant clinical isolates of *Acinetobacter baumannii*. *Infect Drug Resist*. 2019;12:1171-80. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
45. Li S, Duan X, Peng Y, Rui Y. Molecular characteristics of carbapenem-resistant *Acinetobacter* spp. from clinical infection samples and fecal survey samples in Southern China. *BMC Infect Dis*. 2019;19:1-12. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
46. Kanaan MHG, Khalil ZK, Khashan HT, Ghasemian A. Occurrence of virulence factors and carbapenemase genes in *Salmonella enterica* serovar Enteritidis isolated from chicken meat and egg samples in Iraq. *BMC Microbiol*. 2022;22:279. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
47. Bengtsson-Palme J, Kristiansson E, Larsson DGJ. Environmental factors influencing the development and spread of antibiotic resistance. *FEMS Microbiol Rev*. 2018;42(1):fux053 [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
48. Canteón R. Antibiotic resistance genes from the environment: a perspective through newly identified antibiotic resistance mechanisms in the clinical setting. *Clin Microbiol Infect*. 2009;15:20-5. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]

How to Cite:

Rezaeian M, Hoveida L. Prevalence of carbapenem resistance genes in Enterobacteriaceae, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* from meat samples. *Med Lab J*. 2025;19(1):36-41. <http://dx.doi.org/10.29252/mlj.19.1.36>